

Agiflex 20 Ampoules

EAN: 5605481101199 FABRICANTE: DIETMED



AGIFLEX is a food supplement based on Chondractiv™ (containing 55% Type II collagen, 12% Chondroitin Sulfate and 2% Hyaluronic Acid), that is, a patented combination of elements that have a marked chondroprotective action.

DESCRIPTION

What is Agiflex used for?

For the maintenance of cartilage.

Helps lubricate joints and keeps joints flexible.

When is it indicated to take Agiflex?

In the cases of:

- Pain.
- Joint stiffness.
- Cartilage wear.
- Limitation of movement.
- Deformations.
- Crackling.

Ingredients

Water, fructose, grape concentrate, hydrolysed marine collagen*, chondroitin sulphate*, Chondractiv™ (containing 55% Type II Collagen, 12% Chondroitin Sulfate and 2% Hyaluronic Acid), L-ascorbic acid, bioflavonoids, sodium hyaluronate, orange flavour, acidity regulator: citric acid, preservative: potassium sorbate, naringin. *Fish product.

Composition Per 1 ampoule:

- Chondractiv™ 250 mg (containing 55% Type II Collagen, 12% Chondroitin Sulfate and 2% Hyaluronic Acid).



Agiflex 20 Ampoules

EAN: 5605481101199 FABRICANTE: DIETMED



- Total collagen 900 mg.
- Total Chondroitin Sulfate 300 mg.
- Total Hyaluronic Acid 45 mg.
- Vitamin C 120 mg (150% NRV*).
- Bioflavonoids 100 mg.
- Naringin 10 mg.

*Nutrient Reference Value.

Orange flavor.

Instructions for use

1 ampoule per day, preferably before breakfast. It can be dissolved in water or fruit juice.

Contraindications

Not recommended during pregnancy and breastfeeding.

Use should be avoided in asthmatics.

Chondroitin should also be avoided in patients who have had or are experiencing a recurrence of prostate cancer.

Drug Interactions and Others

Minimal side effects have been reported for collagen, mainly some gastrointestinal discomfort, such as a feeling of fullness and unpleasant taste, and headache. Chondroitin sulfate is safe for most people. In some cases it may cause gastric pain and nausea. Other side effects have been reported such as diarrhea, constipation, swollen eyelids, swollen legs, hair loss, and irregular heartbeat. Caution should be taken when combining chondroitin with warfarin (anticoagulant), because it increases the risk of bleeding.

