

Agiflex 40 Capsules

EAN: 5605481107306 FABRICANTE: DIETMED



Vitamin C contributes to the normal formation of collagen for the normal functioning of cartilage.

DESCRIPTION

What is Agiflex used for?

For the maintenance of cartilage.

Helps lubricate joints and keeps joints flexible.

When is it indicated to take Agiflex?

In the cases of:

- Pain.
- Joint stiffness.
- Cartilage wear.
- Limitation of movement.
- Deformations.
- Crackling.

Ingredients

Chondractiv™ (containing 55% Type II Collagen, 12% Chondroitin Sulfate and 2% Hyaluronic Acid), bulking agent: microcrystalline cellulose, marine collagen*, gelatin, L-ascorbic acid, quercetin, sodium hyaluronate, chondroitin sulphate*, naringin, anti-caking agents: magnesium salts of fatty acids, silicon dioxide.

Composition

For 2 capsules: Chondractiv™ 500 mg, Collagen 500 mg, Hyaluronic Acid 50 mg, Chondroitin Sulfate 100 mg, Vitamin C 80 mg (100% VRN*), Quercetin 60 mg, Naringin 20 mg. *Reference Value of Nutrients.

Instructions for use

Prices valid except for changes on our website. (IVA/VAT) included. Doc. Version bf-trf-G-20241116
<https://bio-farma.es/buy/f/dietmed/agiflex-40-capsules>



Agiflex 40 Capsules

EAN: 5605481107306 FABRICANTE: DIETMED



1 capsule 2 times a day, preferably before breakfast and before dinner.

Contraindications

Not recommended during pregnancy and breastfeeding.

Use should be avoided in asthmatics. Chondroitin should also be avoided in patients who have had or are experiencing a recurrence of prostate cancer.

Drug Interactions and Others

Minimal side effects have been reported for collagen, mainly some gastrointestinal discomfort, such as a feeling of fullness and unpleasant taste, and headache. Chondroitin sulfate is safe for most people. In some cases it may cause gastric pain and nausea. Other side effects have been reported such as diarrhea, constipation, swollen eyelids, swollen legs, hair loss, and irregular heartbeat. Caution should be taken when combining chondroitin with warfarin (anticoagulant), because it increases the risk of bleeding.

