

Nodacil Gel 15 ml x 10 Sachets

EAN: 8425652550611 **FABRICANTE: NOVADIET**



Fast and effective relief of symptoms related to heartburn.

DESCRIPTION

What is Nodacil?

It is a dietary supplement based on Magnesium with sweeteners.

What is Nodacil used for?

Nodacil Gel contains magnesium which contributes to electrolyte balance, therefore it is useful for gastric acidity.

How to take Nodacil?

Take one sachet 30-60 minutes after main meals. Recommended daily dose: 1-2 sachets. Shake before opening.

Benefits of taking Nodacil

The sensation of heartburn after meals is caused by the reflux of gastric contents, which are acidic, into the esophagus (an organ not prepared to contain acids).

This occurs due to a failure in the barrier mechanism that separates the esophagus from the stomach. The factors that promote poor functioning of the lower esophageal sphincter are numerous and varied, such as lying down after eating, obesity, tobacco and alcohol, large meals, stress, wearing tight clothing around the waist, some drugs, certain diseases such as hiatal hernia and some foods such as coffee, tea, cola drinks, carbonated drinks, chocolate, mint, citrus fruits, garlic, peppers, etc.

Nodacil Gel is your fast and effective relief from symptoms related to heartburn.

What is Nodacil made of?

Nodacil Gel contains magnesium salts (magnesium carbonate and magnesium trisilicate), calcium salts (calcium carbonate), sodium alginate, pectins, xanthan gum and peppermint essential oil.

Ingredients







Nodacil Gel 15 ml x 10 Sachets

EAN: 8425652550611 FABRICANTE: NOVADIET



Water, Bulking agent (magnesium trisilicate), Magnesium carbonate, Acidity regulator (calcium carbonate), Modified corn starch, Gelling agents (microcrystalline cellulose, carboxymethylcellulose, pectin, xanthan gum, sodium alginate), Peppermint essential oil, Sweeteners (acesulfame K, sucralose), Preservatives (sodium benzoate, potassium sorbate).

Active Ingredients

For 1 sachet:

Magnesium: 125 mg (33% NRV*).

For 2 envelopes:

Magnesium: 250 mg (66% NRV*).

*NRV: Nutrient Reference Values.

Precautions

Do not take this dietary supplement if you are allergic or intolerant to any of its ingredients.

Its use is not recommended during pregnancy and breastfeeding.

Separate taking Nodacil Gel from taking any medication by at least 1 hour.

Take the product with plenty of water to ensure that the substance reaches the stomach. People with swallowing problems are warned of the risk of suffocation. Avoid taking it with medications and other dietary supplements containing fibre.



